

Strengthening our global leadership in treatment of addiction

Morgan Stanley Global Healthcare Conference
September 13th and 14th 2018



Forward Looking Statements

This presentation contains certain statements that are forward-looking and which should be considered, amongst other statutory provisions, in light of the safe harbour provisions of the United States Private Securities Litigation Reform Act of 1995. By their nature, forward-looking statements involve risk and uncertainty as they relate to events or circumstances that may or may not occur in the future. Actual results may differ materially from those expressed or implied in such statements because they relate to future events. Forward-looking statements include, among other things, statements regarding the Indivior Group's financial guidance for 2018 and its medium- and long-term growth outlook, its operational goals, its product development pipeline and statements regarding ongoing litigation.


Various factors may cause differences between Indivior's expectations and actual results, including: factors affecting sales of Indivior Group's products; the outcome of research and development activities; decisions by regulatory authorities regarding the Indivior Group's drug applications; the speed with which regulatory authorizations, pricing approvals and product launches may be achieved; the outcome of post-approval clinical trials; competitive developments; difficulties or delays in manufacturing; the impact of existing and future legislation and regulatory provisions on product exclusivity; trends toward managed care and healthcare cost containment; legislation or regulatory action affecting pharmaceutical product pricing, reimbursement or access; claims and concerns that may arise regarding the safety or efficacy of the Indivior Group's products and product candidates; risks related to legal proceedings, including the ongoing investigative and antitrust litigation matters; the Indivior Group's ability to protect its patents and other intellectual property; the outcome of patent infringement litigation relating to Indivior Group's products, including the ongoing ANDA lawsuits; changes in governmental laws and regulations; issues related to the outsourcing of certain operational and staff functions to third parties; uncertainties related to general economic, political, business, industry, regulatory and market conditions; and the impact of acquisitions, divestitures, restructurings, internal reorganizations, product recalls and withdrawals and other unusual items.



SUBLOCADE™ Update

First SUBLOCADE™ Prescription Journey Timeline – Journey Process Reduction

Original Methodology



Drivers of Journey Time Reduction

- Education of HCP office staff by Field Reimbursement (FRS) team
- Accuracy of patient application
- HUB capacity
- Payor coverage (quantity & quality)
- SP coordination between patient and HCP office



We Continue to See the Number of Patients Receiving Their First Dispense Increase Month Over Month⁽¹⁾

Cumulative Launch to 8/31/18 ⁽¹⁾

77% Payor coverage

9,000+ Unique patient prescription initiations

2,000+ Unique patient injections

Monthly Patient Dispense Conversion data for Q3 not available; updated data will be provided with Q3 results Nov. 1st.

(1) Total includes estimate for patients with HCPs using Specialty Distributors (Buy and Bill)



SUBLOCADE™ (buprenorphine extended-release) Injection KPIs – HCP Data & Patient Treatment Adherence


Cumulative Launch to 8/31/18 (1)

1,700+ HCPs initiated prescription journeys

691 HCPs administered SUBLOCADE™

77 HCPs administered ≥ 5 patients

Treatment Adherence
(All patients with initial injection during March - April)



SUBLOCADE™ Update Conclusion

- Focused on maintaining the momentum on reducing the prescription journey duration and improving HCP trial and success rates
- FY 2018 net revenue expected to be in the range of \$25 million to \$50 million and heavily weighted to Q4 18
- Remain confident in annual peak net revenue goal of \$1 billion+



SUBLOCADE (BUPRENORPHINE EXTENDED-RELEASE) INJECTION FOR SUBCUTANEOUS USE (CIII)

INDICATION AND HIGHLIGHTED SAFETY INFORMATION

INDICATION

SUBLOCADE is indicated for the treatment of moderate to severe opioid use disorder in patients who have initiated treatment with a transmucosal buprenorphine-containing product, followed by dose adjustment for a minimum of 7 days.

SUBLOCADE should be used as part of a complete treatment plan that includes counseling and psychosocial support.

WARNING: RISK OF SERIOUS HARM OR DEATH WITH INTRAVENOUS ADMINISTRATION; SUBLOCADE RISK EVALUATION AND MITIGATION STRATEGY

- **Serious harm or death could result if administered intravenously. SUBLOCADE forms a solid mass upon contact with body fluids and may cause occlusion, local tissue damage, and thrombo-embolic events, including life threatening pulmonary emboli, if administered intravenously.**
- **Because of the risk of serious harm or death that could result from intravenous self-administration, SUBLOCADE is only available through a restricted program called the SUBLOCADE REMS Program. Healthcare settings and pharmacies that order and dispense SUBLOCADE must be certified in this program and comply with the REMS requirements.**

HIGHLIGHTED SAFETY INFORMATION

Prescription use of this product is limited under the Drug Addiction Treatment Act.

CONTRAINDICATIONS

SUBLOCADE should not be administered to patients who have been shown to be hypersensitive to buprenorphine or any component of the ATRIGEL® delivery system

WARNINGS AND PRECAUTIONS

Addiction, Abuse, and Misuse: SUBLOCADE contains buprenorphine, a Schedule III controlled substance that can be abused in a manner similar to other opioids. Monitor patients for conditions indicative of diversion or progression of opioid dependence and addictive behaviors.

Respiratory Depression: Life threatening respiratory depression and death have occurred in association with buprenorphine. Warn patients of the potential danger of self-administration of benzodiazepines or other CNS depressants while under treatment with SUBLOCADE.

Neonatal Opioid Withdrawal Syndrome: Neonatal opioid withdrawal syndrome is an expected and treatable outcome of prolonged use of opioids during pregnancy.

Adrenal Insufficiency: If diagnosed, treat with physiologic replacement of corticosteroids, and wean patient off of the opioid.

Risk of Opioid Withdrawal With Abrupt Discontinuation: If treatment with SUBLOCADE is discontinued, monitor patients for several months for withdrawal and treat appropriately.

Risk of Hepatitis, Hepatic Events: Monitor liver function tests prior to and during treatment.

Risk of Withdrawal in Patients Dependent on Full Agonist Opioids: Verify that patient is clinically stable on transmucosal buprenorphine before injecting SUBLOCADE.

Treatment of Emergent Acute Pain: Treat pain with a non-opioid analgesic whenever possible. If opioid therapy is required, monitor patients closely because higher doses may be required for analgesic effect.

ADVERSE REACTIONS

Adverse reactions commonly associated with SUBLOCADE (in ≥5% of subjects) were constipation, headache, nausea, injection site pruritus, vomiting, increased hepatic enzymes, fatigue, and injection site pain.

For more information about SUBLOCADE, the full Prescribing Information including BOXED WARNING, and Medication Guide visit www.sublocade.com.

